

Evaluations of New Indications

Corticosteroid Aerosols in the Treatment of Asthma

C. S. Ted Tse, Pharm.D., and I. Leonard Bernstein, M.D.

Since the 1950s, corticosteroid aerosols have proved useful in the treatment of asthma. Although their precise mechanism of action is not known, these topical agents have beneficial antiinflammatory and decongestive effects on the bronchial tree in both the allergic and nonallergic forms of this disease. Four of the newer aerosolized steroids — beclomethasone dipropionate, triamcinolone acetonide, flunisolide and budesonide — have been evaluated in clinical trials. The last drug is still investigational. Their side effects are minimal, the major ones being oral candidiasis and dysphonia. They are most effective when used prophylactically and should not be administered during acute asthmatic attacks, as insufficient amounts of drug are inhaled when the airways are constricted. Patients must be instructed in the correct techniques of administering steroid aerosols to ensure optimal therapy. (Pharmacotherapy 1984;4:334-342)

OUTLINE

Structural Activity Relationships

Modes of Action

Beclomethasone Dipropionate

Dosing Frequency

Proper Inhalation Techniques

Use with Other Drugs

Side Effects

Triamcinolone Acetonide

Flunisolide

Pharmacokinetics

Clinical Efficacy

Budesonide

Pharmacokinetics

Comparative Studies

Dosing Frequency

Current Rationale and Recommendations for Corticosteroid Aerosols

Aerosolized preparations of corticosteroids were proposed for the treatment of asthma in the early

1950s, beginning with studies of topical cortisone by Geffard.¹ These studies were performed on the theory that high local concentrations of steroids would be as effective on diseased bronchial mucous membranes as when steroids were applied to the skin or injected into joints. It was also postulated that these locally acting agents might relieve asthma symptoms with fewer side effects than systemically administered steroids. After Geffard reported successful results in 5 patients, other investigators explored the efficacy of hydrocortisone inhaled either as a powder or as a solution.²⁻⁴ They concluded that aerosolized hydrocortisone benefited relatively few patients and gave no evidence that systemic side effects would be diminished.

Dexamethasone phosphate, a water-soluble ester, was the first of the next generation of topical corticosteroids to be studied as an aerosol preparation.⁵ This potent antiinflammatory drug was first investigated in 1962 in the hope that it would gain access to the bronchial tree as far as the terminal respiratory bronchioles and mix with bronchial secretions for better effects. Dexamethasone phosphate was found to be consistently effective in reducing oral steroid requirements and suppressing asthma symptoms in both adults and children.⁶⁻⁸ However, investigators also found that it produced unwanted systemic side effects such as moon facies and decreased urinary cortisol metabolites.⁹⁻¹¹ In addition, Dennis and Ilkin reported that 5 patients developed oral thrush after using dexamethasone aerosol.¹² Although effective in suppressing asthma, this preparation did not seem to have selective topical action on the bronchial mucous membrane, but rather was systemically absorbed in the tracheobronchial tree.¹³

These early clinical experiences revealed that there were no obvious advantages in using hydrocortisone or dexamethasone by the inhalation route.

From the Department of Pharmaceutical Services, University of Chicago Medical Center, and the Department of Internal Medicine, Division of Immunology, University of Cincinnati Medical Center.

Address reprint requests to C. S. Ted Tse, Pharm.D., Department of Pharmaceutical Services, Box 434, University of Chicago Medical Center, Chicago, IL 60637.

BEST AVAILABLE COPY

Therefore it was essential that new inhalation steroid compounds be developed that would provide optimal topical effects but have poor absorption properties.

Structural Activity Relationships

The ideal glucocorticoid for local treatment of asthma would act in the lung but produce a minimum of systemic effects in the therapeutic dose range. To achieve this, the compound should combine properties of selective topical activity with poor absorption or rapid inactivation after absorption (i.e., low bioavailability and short plasma half-life).

For some time it was known that compounds could be synthesized with properties of enhanced glucocorticoid activity and reduced mineralocorticoid effects by modifying the hydrocortisone molecule in positions 1, 2, 6, 9, 16 and 17 (Figure 1). These potent agents (e.g., prednisolone) have poor topical effects, however.¹⁴ To achieve enhanced topical corticosteroid action, some of the hydroxyl groups in the hydrocortisone molecule were substituted with ester acetone groups (Figure 1). Among these potent glucocorticoids with desired topical activity were beclomethasone 17 α , 21-dipropionate and betamethasone 17-valerate (Figure 2). Pharmaceutical chemists also developed compounds, such as triamcinolone acetonide and flunisolide, with desired topical activity by substituting asymmetric 16 α and 17 α acetal groups.¹⁵

It was also demonstrated that the introduction of halogen substituents in the 6 α , 9 α or both positions would increase glucocorticosteroid activity;¹⁶ however, there was no evidence that these halogen substitutions preferentially increased topical activity. Indeed, steroid nucleus fluorination in the 6 α and 9 α positions appeared to potentiate systemic activity more than the topical activity because it decreased systemic metabolism and transcortin binding.¹⁷

Brattsand and co-workers studied the influence of steroid nucleus fluorination on the topical and the systemic activity of 16 α and 17 α acetals and found that to obtain high topical and antiinflammatory activity, optimal fluorination of those groups was more important than it was of the 6 α and 9 α positions.¹⁸ With this theory in mind, chemists developed budesonide, a compound with a high topical:systemic activity ratio (Table 1, Figure 2).

Modes of Action

The mechanism of action of steroids in asthma is not well understood. It is hypothesized that the free steroid molecule diffuses through the cell membrane where it forms an intracellular complex with a specific receptor protein.^{18, 19} This complex is then transported into the cell nucleus where it is bound to specific parts of the chromatin and forms a new type of specific mRNA. This mRNA then determines the DNA and amino acid sequences for the synthesis of new proteins, which are responsible for steroid-specific cellular responses.

HYDROCORTISONE




Figure 1. Structural formula of hydrocortisone. Groups that are essential to antiinflammatory activity are circled.




Figure 2. Structural formulas of 4 topically selective glucocorticosteroids. Budesonide is not yet available in the United States for the treatment of asthma. Flunisolide is also approved as a topical agent for the treatment of allergic rhinitis.

Table 1. Relative Potencies of Topically Selective Glucocorticosteroids in Inhibiting Rat or Mouse Ear Edema Formation and Thymus Involution after Topical Application

Compound	Topical: Antiinflammatory Potency	Systemic Potency	Ratio of Topical to Systemic Potency
Budesonide	1	1	1
Beclomethasone dipropionate	0.4	3.5	0.11
Flunisolide	0.7	12.8	0.05
Triamcinolone acetonide	0.3	5.3	0.05

Adapted from reference 17.

The synthesis of proteins such as lipomodulin and macrocortin in response to steroidal stimulation is believed to inhibit the action of membrane phospholipase A₂ on the arachidonic acid cascade.¹⁵⁻¹⁷ This results in inhibition of phospholipid methylation in the cell membrane and the formation of leukotenes, prostaglandins, thromboxanes and other arachidonic acid metabolites.¹⁸ Consequently, leukotriene-mediated effects such as chemotaxis, histamine release from mast cells and basophils, bronchospasm and inflammatory edema are suppressed.

In theory, the most beneficial effects of steroids in the treatment of asthma are their antiinflammatory and decongestive effects on bronchial mucous membranes. Thus corticosteroids are effective in allergic and nonallergic asthma because the inflammatory process occurs in both conditions.¹⁹ Steroids may stabilize the vascular wall, with reduction of edema and inhibition of migration of inflammatory cells into bronchial tissues.

During allergen challenge, mast cells present in airway lumina release mediators and reduce permeability changes in the usually tight intercellular junctions of the respiratory mucosa, thus facilitating more allergen-mast cell contact and augmenting antigen mediator penetration into the submucosa. Aerosolized steroids are thought to depress these alterations of epithelial and endothelial cell permeability.²⁰

Corticosteroids also possess a conditioning or so-called permissive effect on adrenergic receptors.²¹ Through this effect they may enhance the effects of β -adrenergic agonists in asthma.

Other steroidal effects in controlling asthma include decreased leukocyte function and eosinopenia, which are beneficial in preventing tissue injury.^{22,23}

Beclomethasone Dipropionate

Beclomethasone dipropionate aerosol (BDP) (Beclivent, Vancoril) first became commercially available in the United States as an inhaler in 1976. It has enhanced topical antiinflammatory activity compared to other topical steroids — dexamethasone

phosphate, triamcinolone acetonide and betamethasone valerate — when tested by the vasoconstriction assay method described by McKenzie.²⁴ A dose of 400 μ g of BDP per day is comparable to 5–10 mg of oral prednisone in the control of asthma.²⁵

Also, BDP is shown to be effective in controlling asthmatic symptoms in steroid-dependent adults and children.²⁶⁻²⁸ It has the advantage over oral corticosteroids in that it is relatively free of systemic side effects. It presents no risk to growth and development in pediatric patients.²⁹ Although systemic absorption is insignificant, slight adrenal suppression (based on plasma cortisol determination, urinary free cortisol excretion and the 11-desoxycortisol response to metyrapone) due to inhaled BDP has been reported.³⁰⁻³² In the great majority of patients, this occurred at doses larger than 1600 μ g per day.^{33,34} Klein and colleagues reported no evidence of adrenal suppression in children using 400 μ g of BDP per day.³⁵

The usual daily dose of BDP in controlling asthma in adults is 400 μ g, but some patients may require 800–1000 μ g d. Higher doses may be required particularly in patients whose asthma was previously controlled with 10–15 mg of systemic prednisone daily.³⁶⁻³⁸ Although most patients can eliminate the need for oral steroids by using 400 μ g of BDP per day, those who usually take more than 10 mg of prednisone per day to control their asthma symptoms may benefit from a combination of a small oral dose of prednisone and a higher dose of BDP. Thus BDP achieves the same total corticosteroid effect but with less risk of systemic side effects.

Dosing Frequency

Although the manufacturers recommend using BDP 3–4 times daily in divided doses, less frequent schedules have been reported. Dosing 2 times a day was not significantly different from 4 times daily when both regimens provided the same total daily doses.³⁹ The advantages of the twice-daily regimen were improved patient compliance and possibly fewer local side effects. Since peak effects of steroids subside after 6–8 hours in general, however, it may

be preferable to use a more frequent dosing (i.e., 4 times daily) schedule until a steady state has been reached.³⁶

Proper Inhalation Techniques

Successful corticosteroid aerosol therapy depends on efficient intrapulmonary drug delivery. Many patients do not use the metered-dose inhaler (MDI) skillfully, and as a result treatment may fail by default of the drug delivery system rather than the drug itself. It is therefore very important that patients be instructed about proper inhalation techniques.³⁷ With the MDI held a short distance from the open mouth, the patient should start inspiration from functional residual capacity. Inspiration should be slow and steady. Activation of the inhaler should be after inspiration has begun and the patient should continue inhaling to total lung capacity and hold the breath for a period of 10 seconds. The second actuation of aerosol medication should be done 2 or 3 minutes after the first dose. Patients are instructed about careful mouth washing with tap water after the delivery of inhaled medication.

The use of a spacer fitted into the MDI has been recommended in asthmatic children and in patients who use their conventional MDI ineffectively. Spacers have been shown to decrease oropharyngeal deposition of inhaled aerosols and to increase intrapulmonary deposition of aerosolized materials.³⁸ The frequency of oropharyngeal complications and the need for antifungal therapy are greatly reduced with the use of spacers. They would be particularly useful for selected patients whose response to inhaled steroid is compromised by dose-limiting oropharyngeal complications and to those who need a greater antiasthmatic effect.³⁹

Use with Other Drugs

Although BDP can be used concurrently with other antiasthmatic drugs, it should be reserved for patients who fail to respond to conventional asthma therapy (Figure 3). Most patients with mild asthma can be treated successfully with bronchodilator drugs (beta agonists, theophylline) and cromolyn sodium. In severe disease or when asthma symptoms become chronic and labile, corticosteroids should be used. Like cromolyn sodium, BDP has no effect during an acute attack of asthma, and should not be used for this purpose.

In general, more than 50% of steroid-dependent patients with asthma can be maintained free of oral steroids most of the time while taking BDP.²⁶ Patients previously unable to convert to alternate-day prednisone can do so during BDP therapy.³⁹ Care must be taken, however, when switching from systemic to aerosol steroids. Oral prednisone must be tapered slowly because sudden discontinuation could cause uncomfortable systemic symptoms or even precipitate adrenal failure. During periods of stress, upper respiratory infections or exacerbations




Figure 3. Steps in treating chronic asthma. Adapted from reference 28.

of severe asthma, it is crucial that generous amounts of oral steroids be added to the aerosol regimen until the acute crisis has subsided.

Side Effects

The most widely reported side effect of BDP is oropharyngeal candidiasis (oral thrush). It is dose-related and causes symptoms in about 5% of patients.^{23,26} It is relatively trivial from the clinical point of view and usually responds to a single course of nystatin. Patients need to be reminded about rinsing their mouths with water or mouthwash after each inhalation dose. Dysphonia affects up to 50% of users and occasionally can be severe and persistent. Toogood and associates reported the advantages of using two spacers in aerosolized corticosteroid delivery to reduce oropharyngeal candidiasis and double antiasthmatic potency.³⁸

Other associated symptoms such as exacerbation of eczema or rhinitis may occur when a patient whose symptoms were previously suppressed by oral steroids is changing from systemic steroids to aerosol BDP.

Mild suppression of adrenal function can occur if more than 1600 $\mu\text{g/d}$ of BDP is used.²⁹ The drug is reported to be safe during pregnancy when recommended doses are used.⁴⁰ When flunisolide nasal

solution and BDP were used together, two independent groups reported that there were no significant additive effects on plasma cortisol levels or on frequency of oral candidiasis.⁴¹⁻⁴² Although a decrease of serum IgG may occur with the use of BDP, there is no reported increase in the occurrence of bacterial infections.⁴³

Triamcinolone Acetonide

Triamcinolone acetonide (Azmecort, Rorer) is a nonpolar, water-insoluble fluorinated steroid that has recently become available for commercial use in the United States. It comes in an aerosolized form in an MDI attached to a barrel-shaped spacer. As previously discussed, initial trials with water-soluble corticosteroids, such as dexamethasone phosphate, did not show any significant advantage over oral steroids,^{3,11-13} however, development of nonpolar, water-insoluble preparations of topical steroids would be expected to have distinct advantages over water-soluble ones in that they have less potential for systemic absorption and thus have a lower risk of side effects.⁴⁴⁻⁴⁵ Topical antiinflammatory potency of triamcinolone acetonide (Figure 2) is less than that of BDP.⁴⁶⁻⁴⁷

The manufacturer of triamcinolone acetonide aerosol (TAA) designed a special spacer delivery system that minimizes deposition of drug in the oral cavity and increases efficiency of drug access to the lungs. One puff of this aerosol releases 200 μ g of TAA, 100 μ g of which are delivered from the unit to the airways as a fine suspension. The recommended daily dose is 2 inhalations 3-4 times a day. In delivered doses ranging from 400-2000 μ g d, TAA has proved to be effective therapy for asthma in both long-term and short-term studies.⁴⁸⁻⁵³

The efficacy and safety of TAA in steroid-dependent asthmatic patients were studied by Bernstein et al in a multicenter, short-term controlled and long-term open study.⁵² Patients treated with TAA in doses of 800-1600 μ g d showed highly significant improvement from baseline in pulmonary function tests and in asthmatic symptoms, whereas no significant improvement was observed in patients treated with placebo aerosol. Mean changes in plasma cortisol level were not statistically significant after 1 year. The only significant side effects reported in this study were hoarseness and sore throat.

Three long-term studies (3-26 months) investigated the effects of TAA in reducing oral steroid requirements in heavily steroid-dependent patients.⁵⁴⁻⁵⁶ Forty percent to 70% of patients discontinued the use of oral steroids and were maintained solely on TAA. Several patients developed symptoms of adrenal insufficiency while tapering prednisone, but some demonstrated disappearance of cushingoid features as well as improvement in asthmatic symptoms.⁴⁹ Minimal side effects of mild hoarseness and periodic loss of voice in 1 patient were observed.⁴⁹ In four short-term studies of 4-12 weeks' duration, TAA was studied for its effectiveness in controlling

asthma symptoms and the feasibility of eliminating or reducing oral steroids.^{46, 50, 51, 53} In doses ranging from 400-1400 μ g d of TAA, asthma remained under satisfactory control, pulmonary function improved significantly, oral steroids were either reduced or totally withdrawn and plasma cortisol levels increased in three out of four studies. Oral candidiasis was reported in 2 patients in one of these studies.⁵⁰ Treatment of asthma in children with TAA was also reported to be successful, with no evidence of adrenal suppression.⁵⁴ Side effects are minimal, since there is a low frequency (2.5%) of hoarseness and oral candidiasis.⁵⁵

Flunisolide

Flunisolide is another synthetic corticosteroid with potent antiinflammatory activity. The aerosolized form has been studied for the treatment of asthma. Its systemic potency is equivalent to that of triamcinolone and is about 5 times greater than that of cortisol.⁵⁶ It is fluorinated in the 6 position and is polar because of cyclized acetonide in the 16 and 17 positions (Figure 2). Flunisolide aerosol was recently approved by the Food and Drug Administration and will be marketed under the trade name of Aerobid by Key Pharmaceuticals.

Pharmacokinetics

Pharmacokinetic properties of flunisolide have been determined in healthy volunteers. The drug is absorbed through pulmonary membranes, and buccal and intestinal mucosae.⁵⁷ Systemic availability of a single inhaled dose (1 mg) of flunisolide was 30-40%. Peak plasma concentration was achieved in 2 minutes and maintained at approximately this level for about an hour. The elimination rate of flunisolide was similar after administration by intravenous, oral and inhalation routes with the terminal elimination half-life being 1.82 ± 0.42 hours. Rapid degradation of systemic flunisolide occurred by extensive first-pass metabolism to the 6 β -OH metabolite and other water-soluble conjugates, which are relatively inactive.⁵⁸

The rapid metabolism of flunisolide apparently accounts for its low systemic toxicity. The drug can be administered in therapeutically effective doses for several months in adults and children without their showing significant systemic effect (see section below).

The relative potencies of intravenous/oral and intravenous/inhalation routes for flunisolide are 6:1 and 3:1 respectively, as assessed by its suppression of eosinophils.⁵⁷

Clinical Efficacy

Clinical investigators have reported effective results of flunisolide aerosol for asthma in both blinded and open studies of adults and children.⁵⁹⁻⁶⁴ In three separate double-blind trials conducted in adults with chronic asthma, flunisolide aerosol ranging from

0.8–2 mg daily applied in 2 separate doses was compared with placebo for 3 weeks to over 1 year. The investigators found statistically significant differences in favor of flunisolide in major therapeutic responses: improvement in spirometric function and daily symptom scores, reduction or elimination of oral steroids and reduction of nonsteroidal asthma medications.^{62–64} Some patients developed oral candidiasis when flunisolide was used for over a year, but there were no major adverse clinical reactions or laboratory abnormalities.

In another double-blind multicenter study in adults with steroid-dependent asthma, 40 patients received 1 mg/d flunisolide aerosol and 33 received placebo for 16 weeks.⁶⁵ There was a significant difference in the decreases in median oral prednisone dosage between the study groups: 74.4% in the flunisolide group compared to only 4.2% in the placebo group. Complete withdrawal of oral steroids was achieved in 27.5% of flunisolide-treated patients and in 12.1% of patients receiving placebo. Reduction in the frequency of asthmatic attacks also favored those treated with flunisolide. Although there were no statistically significant differences in asthma severity between the groups prior to the study, a decrease in severity occurred during the study in both groups, but was greater in the patients receiving flunisolide.⁶⁵ These patients were also associated with an increase in plasma cortisol level (43%), but no change was observed in the placebo group. The authors concluded that flunisolide aerosol (1 mg/d) provided superior symptomatic control and replaced an average of 9 mg/d of oral prednisone without causing serious adverse local or systemic effects.

Webb et al obtained similar results.⁶⁶ In a 3-month study, they studied the efficacy of flunisolide aerosol in 16 steroid-dependent and 13 steroid-independent patients with asthma. Each patient received 2 mg of flunisolide daily by aerosol (four 250- μ g puffs twice daily). Adrenal function was monitored with repeated measurements of plasma cortisol levels and metyrapone tests. The steroid-dependent patients demonstrated a significant decrease in use of prednisone (61%) and adrenergic aerosol, improved pulmonary function and a significant increase in morning plasma cortisol levels (33%). Steroid-independent patients had significant improvement in forced expiratory volume in 1 second (FEV₁) and a decrease in the use of theophylline and adrenergic aerosols. They had no change in morning cortisol levels, however. Side effects, which were minimal in both groups, included throat irritation, abdominal bloating and the appearance of nasal polyps and rhinitis in some steroid-dependent patients after discontinuing systemic steroid therapy.⁶⁶

Efficacy of flunisolide aerosol was also addressed in pediatric populations.^{61, 66, 67} Shapiro and associates studied 32 steroid-dependent asthmatic children for a period of 12 weeks using a double-blind, prerandomized method.^{61, 67} The drug-treated group received 0.5 mg of flunisolide aerosol twice a day. All

patients receiving flunisolide had improved asthma control and a significant decrease in oral steroid dosage. Only half of the placebo-treated control group had a decline in steroid requirements. Pulmonary function tests and adrenal function remained stable in both groups. Patients in both groups developed some pharyngeal candidiasis, but this could have been related to previous steroid use in some of them.

Meltzer and others conducted another short-term, double-blind, placebo-controlled study lasting 8 weeks. The 46 steroid-independent asthmatic children⁶⁸ received either 0.5 mg flunisolide aerosol or placebo twice a day. Effectiveness was evaluated by daily symptom scores, pulmonary function and physical examination. Most symptom scores (severity of wheezing, chest tightness, shortness of breath and frequency and severity of asthma attacks) were significantly better in those receiving flunisolide than in those receiving placebo. Pulmonary function showed improvement in the flunisolide-treated group and deterioration in the placebo group, but these differences were not statistically significant. No serious adverse effects were reported. No patient developed thrush or evidence of adrenal suppression.

Gale and co-workers evaluated the effects of concurrent administration of flunisolide and beclomethasone in patients with both rhinitis and asthma.⁶⁴ Flunisolide nasal solution was added to either BDP bronchial aerosol or flunisolide bronchial aerosol to study the possible cumulative effects of topical corticosteroids on adrenal function. The investigators also evaluated the efficacy of these combinations in controlling symptoms associated with rhinitis and asthma for 1 month. Patient and physician assessments revealed no significant differences between the combinations in efficacy, adverse effects or effect on adrenal function.

The frequency of oropharyngeal candidiasis in patients using flunisolide aerosol and BDP was studied by Spector and colleagues.⁶⁸ Symptomatic thrush was slightly more common ($p = 0.03$) in patients treated with BDP (200 μ g 4 times a day) compared to those taking flunisolide (500 μ g twice a day). Patients with pretrial throat cultures that were positive for candida organisms had significantly greater frequency of clinical thrush than those with negative cultures. Positive pretrial throat cultures provided a good indication of patients at higher risk of developing thrush; they might benefit from more frequent and prolonged mouthwashes after using corticosteroid aerosols.

Budesonide

Budesonide is an investigational 16 α , 17 α -acetal corticosteroid that is not halogenated (Figure 2). It has a high degree of topical antiinflammatory activity and low systemic potency.⁶⁹ After topical application, it was 2 times as potent as beclomethasone and 3 times as potent as flunisolide in inducing vasoconstriction.⁶⁹ It lacks the halogen substitution that is present in flunisolide, triamcinolone and beclometh-

asone. Since this substitution decreases the rate of biotransformation, budesonide is rapidly metabolized¹⁵ (Table 2).

Pharmacokinetics

Within seconds after the drug was administered by bronchial inhalation in healthy volunteers, unchanged budesonide was detected in blood plasma, indicating that the drug is absorbed intact through the respiratory tract.⁷² This initially high plasma concentration also indicated minimal metabolism of drug in the lung. Plasma half-life of unchanged budesonide was estimated to be 2.0 ± 0.2 hours, a value similar to that found after intravenous injection (2.8 ± 1.1 hr).

Human pharmacokinetic studies showed that budesonide is readily biotransformed in the liver by oxidative and reductive biotransformation, but not in the lung or skin.⁶⁹ The systemic availability after oral administration was calculated to be $10.7 \pm 4.3\%$. Extensive first-pass metabolism occurred as the drug biotransformed rapidly when incubated with human liver.⁷³ In vitro biotransformation studies in human liver isolated two major metabolites, 6 β -hydroxybudesonide and 16 α -hydroxyprednisolone.⁷⁴ They are virtually inactive and were 1/10 to 1/100 weaker than the parent compound. Plasma protein binding was around 88%. Recovery studies in humans after inhalation of ³H budesonide showed that most of the radioactivity was excreted in the urine (32%) and feces (15%).⁷⁵ About 40% of the administered dose was deposited in the inhaler and about 5% was deposited in the oral cavity.

Pharmacologic studies in rats showed that the ratio of topical to systemic effects of budesonide was 10–20 times better than that of BDP and TAA⁷⁶ (Table 1). Such improvement makes budesonide a promising alternative for aerosol treatment of asthma.

Eliu-Micallef et al administered 1 mg of budesonide by inhalation to 12 patients with chronic bronchial asthma and studied responses over time.⁷⁷ Two hours after drug inhalation, a statistically significant increase in peak expiratory flow (PEF) occurred. Peak effect occurred between 6 and 7 hours after budesonide inhalation and the change in bronchial function was still significant for up to 12 hours.

Comparative Studies

The clinical efficacy of aerosol budesonide in bronchial asthma has been compared satisfactorily with that of aerosol BDP, subcutaneous terbutaline and oral prednisolone.^{71–75}

In open short-term, crossover trials, budesonide was compared to BDP in 27 steroid-dependent asthmatics.⁷² Both drugs were given in the same dosage of 200 μ g 4 times a day for 2 weeks and no significant differences were noted between them. The report did not mention any side effects.

Willey et al performed a well-designed, double-blind crossover trial comparing 100 μ g of BDP 4

Table 2. Plasma Half-lives of Various Corticosteroids

Corticosteroid	Plasma Half-life (min)
Cortisone	90
Cortisol	90
Flunisolide	100
Budesonide	150
Prednisolone	200
Methylprednisolone	200
Triamcinolone	200
Dexamethasone	300
Betamethasone	300
Beciomethasone ciproionate	900 ^a

Adapted from references 80 and 81.

^aFrom reference 81.

times a day and 200 μ g of budesonide 2 times a day in 30 chronic asthmatics.⁷² Each treatment regimen was studied for 4 weeks. Assessments consisted of daily scores of asthma severity and morning and evening PEF values. Every 2 weeks the authors measured forced vital capacity (FVC) and FEV₁. There were no significant differences in PEF values during the study period. Although FEV₁ values were slightly higher after budesonide therapy ($p < 0.05$), FVC values showed no significant differences. During the 8 weeks, there were no clinically significant systemic side effects or cases of oropharyngeal candidiasis. Similarly, symptom scores between the groups showed no significant differences. Thus budesonide 200 μ g twice a day was at least as effective as BDP 100 μ g 4 times a day in controlling asthma symptoms.

Dahl and Jonansson performed double-blind, placebo-controlled study in 21 asthmatics to investigate the clinical effects of inhaled budesonide and subcutaneous terbutaline given separately or simultaneously.⁷⁸ Terbutaline produced a faster onset of bronchodilation than budesonide (1 hr vs 4 hrs) as indicated by changes of PEF. The peak increase in PEF after either drug favored terbutaline. The two drugs given simultaneously did not show potentiation or additive effect.

Contrary to the above study, Henriksen and Dahl found that when inhaled terbutaline 32.5 μ g and inhaled budesonide 400 μ g were given together to 14 children with exercise-induced asthma, the drug combination exerted an additive effect on improving pulmonary function.⁷⁹ The authors also concluded that 1–4 weeks of treatment with inhaled budesonide decreased the severity of asthma in these children. No side effects were reported.

Several short-term trials studied the clinical effects after oral prednisolone and inhaled budesonide in asthma patients.^{71–75} The investigators found that 400 μ g of budesonide had the same effect on PEF as did 10 mg of prednisolone, whereas 800 μ g of bude-

sonide was as potent as 20 mg of prednisolone.⁷⁵ However, the authors did not specify the statistical methods used to evaluate the data. Another double-blind trial administered 40 mg prednisolone orally, placebo orally and 1 mg of budesonide by inhalation. Oral prednisolone produced a statistically significantly greater effect on PEF from the eighth hour after drug administration onward.⁷¹ The peak effect occurred between 6 and 7 hours after budesonide inhalation and about 9 hours after prednisolone. Oral prednisolone 40 mg also produced a significantly greater maximal response in PEF, but the response peaked 3 hours later than budesonide.

Budesonide exerted a dose-dependent suppression of blood eosinophils and plasma cortisol levels.^{76,77} This change is significant if budesonide is used in doses of 1600 µg/d.⁷⁷ The drug has less systemic effects on eosinophils and plasma cortisol levels when compared with BDP.^{69,77}

Dosing Frequency

Several investigators studied the influence of dosing frequency on the efficacy of aerosol budesonide in asthmatic patients. The results were contradictory.⁷²⁻⁷⁴ Data from Toogood et al showed that the same daily dose given in different dosing schedules did not affect the drug's systemic potency but it did affect its antiasthmatic potency.⁷² Twice-daily dosing was associated with reduction in drug potency as indicated by the drop in daily PEF and increase in severity of symptoms.⁷² This agreed with the findings of Dahl and Jonansson⁷³ but contrasted with those of Stiska et al.⁷⁴ who claimed that reduction of the frequency of inhaled budesonide did not reduce the effect of the treatment. As asthma became more severe, however, dosing frequency became a major response-limiting factor, and could not be offset despite a fourfold increase in daily dosage.⁷³⁻⁷⁴ Less frequent dosing decreased the prevalence of oropharyngeal candidiasis.⁷² Overall, administration 4 times a day provided the best risk-benefit relationship.

Current Rationale and Recommendations for Corticosteroid Aerosols

Many controlled studies have shown that corticosteroid aerosols such as beclomethasone dipropionate, triamcinolone acetonide, flunisolide and budesonide are highly effective in the control of asthma. All of these agents have high topical activity, low systemic activity and rapid hepatic metabolism (Table 1 and 2). Side effects are minimal, the major ones being oral thrush and hoarseness. Used within the normal range, these drugs rarely suppress adrenal function and are safe for both adults and children.

Theophylline derivatives, beta-agonistic bronchodilators and cromolyn sodium should all be tried before either aerosol or systemic steroids are administered.^{82,83} Some patients may require the use of bronchodilator aerosol 10-15 minutes before inhal-

ing a steroid aerosol, thus allowing the aerosol to penetrate deeper into the peripheral airways. Short courses of systemic steroids may be necessary during periods of stress or reduced airway patency. Immunotherapy should not be stopped during treatment with oral or inhaled steroids.

Patients should be reminded that aerosol steroids are for prophylactic use only and should be used regularly to prevent asthma. These preparations should not be used during an acute attack of asthma because a smaller fraction of the dose will be inhaled due to airway obstruction.

Finally, all patients taking steroid aerosols should be observed and instructed by both physicians and pharmacists about the proper use of the inhaler spacer in order to experience optimal therapy.

References

- Gelfand ML. Administration of cortisone by the aerosol method in the treatment of bronchial asthma. *N Engl J Med* 1951;245:293-4.
- Foulds WS, Greaves DP, Herxheimer H, Kingdom LG. Hydrocortisone in treatment of allergic conjunctivitis, allergic rhinitis and bronchial asthma. *Lancet* 1955;1:234-5.
- Brockbank W, Pengelly CDR. Chronic asthma treated with powder inhalation of hydrocortisone and prednisone. *Lancet* 1958;1:187-2.
- Heim WH, Heyworth F. Bronchial asthma and chronic bronchitis treated with hydrocortisone acetate inhalation. *Br Med J* 1958;2:765-8.
- Herxheimer H, McAllen MK, Williams DA. Local treatment of bronchial asthma with hydrocortisone powder. *Br Med J* 1958;2:762-5.
- Smith JM. Hydrocortisone hemisuccinate by inhalation in children with asthma. *Lancet* 1958;2:1248-9.
- Bickerman HA, Itkin SE. Aerosol steroid therapy and chronic bronchial asthma. *JAMA* 1963;184:533-8.
- Crepea SB. Inhalation corticosteroid (dexamethasone PO₁) management of chronically asthmatic children. *J Allergy* 1963;34:119-26.
- Arbesman CE, Bonstein HS, Reisman RE. Dexamethasone aerosol therapy for bronchial asthma. *J Allergy* 1963;34:354-61.
- Novay HS, Beall G. Aerosolized steroids and induced Cushing's syndrome. *Arch Intern Med* 1965;115:602-5.
- Siegel SC, Heimlich EM, Richards W, Kelly VC. Adrenal function in allergy. IV. Effect of dexamethasone aerosols in asthmatic children. *Pediatrics* 1964;33:245-50.
- Dennis M, Itkin IH. Effectiveness and complications of aerosol dexamethasone phosphate in severe asthma. *J Allergy* 1964;35:70-5.
- Toogood JH, Letcoe NM. Dexamethasone aerosol for the treatment of steroid-dependent bronchial asthmatic patients. *J Allergy* 1965;35:321-32.
- Franklin W, Lowell FC, Michaelson AL, Schiller IW. Aerosolized steroids in bronchial asthma. *J Allergy* 1958;29:214-17.
- Sarett LH, Patchett AT. The effects of structural alteration on the antiinflammatory properties of hydrocortisone. *Prog Drug Res* 1963;5:11-153.
- Popper TL, Watnick AS. Anti-inflammatory steroids. In: Scheraga RA, Whitehouse MW, eds. *Anti-inflammatory agents*. Vol 1. New York: Academic Press, 1974:245-34.
- Brattsand R, Thalen A, Roempke K, Kallstrom L, Grusstad E. Development of new glucocorticoids with a very high ratio between local and systemic activities. *Eur J Respir Dis* 1982;63(suppl 122):62-73.
- Baxter JD, Funder JW. Hormone receptors. *N Engl J Med* 1979;301:1149-61.
- Baxter JD, Forsham PH. Tissue effects of glucocorticoids. *Am J Med* 1972;53:573-89.
- Blackwell GJ, Carnuccio R, Dirosa M, Flower RJ, Parente L, Perissic P. Macro cortin: a polypeptide causing antihypertensive effect of glucocorticoids. *Nature* 1980;287:147-9.
- Hirata F, Schiffman E, Venkatasubramanian K, Solomon D, Axelrod J. A phospholipase A₂ inhibitory protein in rabbit neutrophils induced by glucocorticoids. *Proc Natl Acad Sci USA* 1980;77:2533-6.
- Kay AB. Basic mechanisms in allergic asthma. *Eur J Respir Dis* 1982;63(suppl 122):9-16.
- Svedmyr N. Mode of action of corticosteroids. In: Mygind N, Clark TJH, eds. *Topical steroid treatment for asthma and rhinitis*. London: Bailliere Tindall, 1980:1-11.
- Webb DR. Steroids in allergic disease. *Med Clin North Am* 1981;65:1073-81.

25. Butterworth AE, Wasson DL, Gleich GJ et al. Damage to schistosoma induced by eosinophil major basic protein. *J Immunol* 1979; 122:221-9.
26. Svedmyr N. Effect of glucocorticoids on the airways. *Eur J Respir Dis* 1982;63(suppl 122):48-53.
27. McKenzie AW. Percutaneous Absorption of Steroids. *Arch Dermatol* 1962;86:611-4.
28. Clark TJH. Beclomethasone dipropionate treatment of asthma in adults. In: Mygind N, Clark TJH, eds. Topical steroid treatment for asthma and rhinitis. London: Bailliere Tindall, 1980:94-106.
29. Toogood JH, Lefcoe NM, Haines DSM et al. Minimum dose requirements of steroid-dependent asthmatic patients for aerosol beclomethasone and oral prednisone. *J Allergy Clin Immunol* 1978;61:355-64.
30. Godfrey S, Balfour-Lynn L, Tooley M. A three- to five-year follow-up of the use of aerosol steroid, beclomethasone dipropionate, in childhood asthma. *J Allergy Clin Immunol* 1978;62:335-9.
31. Wyatt R, Waschek J, Weinberger M, Sherman B. Effects of inhaled beclomethasone dipropionate and alternate-day prednisone on pituitary-adrenal function in children with chronic asthma. *N Engl J Med* 1978;299:1387-92.
32. Smith MJ, Hodson ME. High-dose beclomethasone inhaler in the treatment of asthma. *Lancet* 1983;1:265-9.
33. Klein R, Waldman D, Kershner H et al. Treatment of chronic childhood asthma with beclomethasone dipropionate aerosol. A double-blind crossover trial in nonsteroid-dependent patients. *Pediatrics* 1977;60:7-13.
34. Costello JF, Clark TJH. Response of patients receiving high-dose beclomethasone dipropionate. *Thorax* 1974;29:571-3.
35. Anonymous. Double-blind trial comparing two dosage schedules of beclomethasone dipropionate aerosol in the treatment of chronic bronchial asthma. preliminary report of the Brompton Hospital Medical Research Council Collaborative Trial. *Lancet* 1974;2:303-7.
36. Munch EP, Taudorf E, Weeke B. Dose frequency in the treatment of asthmatics with inhaled topical steroid. *Eur J Respir Dis* 1982;63(suppl 122):143-53.
37. Harper TB, Strunk RC. Techniques of administration of metered-dose aerosolized drugs in asthmatic children. *Am J Dis Child* 1981;135:218-21.
38. Toogood JH, Baskerville J, Jennings B, Lefcoe NM, Johansson SA. Use of spacers to facilitate inhaled corticosteroid treatment of asthma. *Am Rev Respir Dis* 1984;129:723-9.
39. Toogood JH, Jennings B, Greenway RW, Chuang L. Candidiasis and dysphonia complicating beclomethasone treatment of asthma. *J Allergy Clin Immunol* 1980;65:145-53.
40. Greenberger PA, Patterson R. Beclomethasone dipropionate for severe asthma during pregnancy. *Ann Intern Med* 1983;98:478-80.
41. Rusnak SL. Concurrent administration of flunisolide nasal solution with beclomethasone dipropionate bronchial aerosol in patients with both rhinitis and asthma. *Ann Allergy* 1981;47:320-4.
42. Gale AE, Harding P, Solomon E. Flunisolide intranasal solution compared with intrabronchial steroid in adults with both bronchial asthma and perennial rhinitis. *Ann Allergy* 1981;46:268-72.
43. De Cotis BA, Settipane GA. The effect of inhaled beclomethasone on serum immunoglobulins. *WESA Proceedings* 1981;2:153. Cited in: Spector SL. The use of inhaled corticosteroid aerosols in the treatment of asthma. *Modest Formulary* 1983;19:421-6.
44. Morrow Brown H, Storey G, George WHS. Beclomethasone dipropionate: a new steroid aerosol for the treatment of allergic asthma. *Br Med J* 1972;1:335-90.
45. Clark TJH. Effects of beclomethasone dipropionate delivered by aerosol in patients with asthma. *Lancet* 1972;1:1361-4.
46. Falliers CJ. Triamcinolone acetonide aerosols for asthma. I. Effective replacement of systemic corticosteroid therapy. *J Allergy Clin Immunol* 1976;57:1-11.
47. Kriz RJ, Chmelik F, dePico G, Reed CE. A one-year trial of triamcinolone acetonide aerosol in severe steroid-dependent asthma. *Chest* 1977;72:36-44.
48. Williams MH. Treatment of asthma with triamcinolone acetonide aerosol. *Chest* 1975;68:765-8.
49. Williams MH, Kane C, Shim CS. Treatment of asthma with triamcinolone acetonide delivered by aerosol. *Am Rev Respir Dis* 1974;109:538-43.
50. Grieco MH, Dwek J, Larsen K, Rammohan G. Clinical effect of aerosol triamcinolone acetonide in bronchial asthma. *Arch Intern Med* 1978;138:1337-41.
51. Kriz RJ, Chmelik F, dePico G, Reed CE. A short-term double-blind trial of aerosol triamcinolone acetonide in steroid-dependent patients with severe asthma. *Chest* 1976;69:455-60.
52. Bernstein IL, Chervinsky P, Falliers CJ. Efficacy and safety of triamcinolone acetonide aerosol in chronic asthma. Results of a multicenter, short-term controlled and long-term open study. *Chest* 1982;81:20-6.
53. Chervinsky P. Treatment of steroid-dependent asthma with triamcinolone acetonide aerosol. *Ann Allergy* 1977;38:192-7.
54. Sly RM, Imseis M, Frazer M, Joseph F. Treatment of asthma in children with triamcinolone acetonide aerosol. *J Allergy Clin Immunol* 1978;62:76-82.
55. Chervinsky P, Petraco AJ. Incidence of oral candidiasis during therapy with triamcinolone acetonide aerosol. *Ann Allergy* 1979;43:80-3.
56. Webb DR, Mullarkey MF, Freeman ML. Flunisolide in chronic bronchial asthma. *Ann Allergy* 1979;42:80-2.
57. Chaplin MD, Cooper WC, Segre EJ, Oren J, Jones RE, Nerenberg C. Correlation of flunisolide plasma levels to eosinopenic response in humans. *J Allergy Clin Immunol* 1980;65:445-53.
58. Chaplin MD, Rooks W, Swenson EW, Cooper WC, Nerenberg C, Chu NI. Flunisolide metabolism and dynamics of a metabolite. *Clin Pharmacol Ther* 1980;27:402-13.
59. Schulz JL, Johnson JD, Freedman SO. Double-blind trial comparing flunisolide and placebo for the treatment of perennial rhinitis. *Clin Allergy* 1978;8:313-20.
60. Sahay JN, Chatterjee SS, Engler C. Flunisolide — a new intranasal steroid for the treatment of allergic rhinitis. *Clin Allergy* 1979;9:17-24.
61. Shapiro GG, Izu AE, Furukawa CT, Pierson WE, Bierman CW. Short-term double-blind evaluation of flunisolide aerosol for steroid-dependent asthmatic children and adolescents. *Chest* 1981;80:671-5.
62. Lowell FC, Ohman JL, Williams M. Double-blind trial of inhaled flunisolide in bronchial asthma. *J Allergy Clin Immunol* 1976;57:257.
63. Zeitz H, Luskin A, Kentor P. Efficacy and safety of flunisolide aerosol in bronchial asthma. *Ann Allergy* 1977;39:70.
64. Spangler OL, Bloom FL, Brestel EP, Wittig HJ. A one-year trial of aerosolized flunisolide in severe steroid-dependent asthmatics. *Ann Allergy* 1977;39:70.
65. Slavin RG, Izu AE, Bernstein IL et al. Multicenter study of flunisolide aerosol in adult patients with steroid-dependent asthma. *J Allergy Clin Immunol* 1980;66:379-85.
66. Meltzer EO, Kemp JP, Orgel HA, Izu AE. Flunisolide aerosol for treatment of severe, chronic asthma in steroid-independent children. *Pediatrics* 1982;69:340-5.
67. Shapiro GG, Furukawa CT, Pierson WE, Bierman CW. Double-blind controlled study of flunisolide aerosol in steroid-dependent asthmatic children. *J Allergy Clin Immunol* 1979;63:163.
68. Spector SL, Wangaard C, Bardana EJ. The use of cultures and immunologic procedures to predict oropharyngeal candidiasis in patients on steroid aerosols. *Clin Allergy* 1982;12:259-78.
69. Johansson SA, Anderson KE, Brattstad R, Grusvstad E, Hedner P. Topical and systemic glucocorticoid potencies of budesonide, beclomethasone dipropionate and prednisone in man. *Eur J Respir Dis* 1982;63(suppl 122):71-82.
70. Ryrfeldt A, Anderson P, Edsbacker S, Tonnesson M, Davis O, Pauwels R. Pharmacokinetics and metabolism of budesonide, a selective glucocorticoid. *Eur J Respir Dis* 1982;63(suppl 122):86-95.
71. Ellul-Micallef R, Hansson E, Johansson SA. Budesonide: a new corticosteroid in bronchial asthma. *Eur J Respir Dis* 1980;61:167-72.
72. Stiksa G, Glennow C, Johannesson N. An open cross-over trial with budesonide and beclomethasone dipropionate in patients with bronchial asthma. *Eur J Respir Dis* 1982;63(suppl 122):266-7.
73. Dahl R, Johansson SA. Effect on lung function of budesonide by inhalation, terbutaline SC and placebo given simultaneously or as single treatments. *Eur J Respir Dis* 1982;63(suppl 122):132-7.
74. Willey RF, Godden OJ, Carmichael J, Preston P, Frame M, Crompton GK. Comparison of twice-daily administration of a new corticosteroid budesonide with beclomethasone dipropionate four times daily in the treatment of chronic asthma. *Br J Dis Chest* 1982;76:61-4.
75. Rosenhall L, Lundqvist G, Adenroth E, Glennow C. Comparison between inhaled and oral corticosteroids in patients with chronic asthma. *Eur J Respir Dis* 1982;63(suppl 122):154-62.
76. Henriksen JM, Dahl R. Effects of inhaled budesonide alone and in combination with low-dose terbutaline in children with exercise-induced asthma. *Am Rev Respir Dis* 1983;128:993-7.
77. Toogood JH, Baskerville JC, Jennings B, Lefcoe NM, Johansson SA. Influence of dosing frequency and schedule on the response of chronic asthmatics to the aerosol, budesonide. *J Allergy Clin Immunol* 1982;70:288-98.
78. Dahl R, Johansson SA. Clinical effect of bid and qid administration of inhaled budesonide. A double-blind controlled study. *Eur J Respir Dis* 1982;63(suppl 122):268-9.
79. Toogood JH. Concentrated aerosol formulations in asthma. *Lancet* 1983;1:790-1.
80. Pauwels R, Van Den Straeten M. Human pharmacokinetics of glucocorticoids. *Eur J Respir Dis* 1982;63(suppl 122):83-5.
81. Schering Laboratories, Kenilworth, NJ. 1981.
82. Bernstein IL, Johnson CL, Tse CST. Therapy with cromolyn sodium. *Ann Intern Med* 1978;89:228-33.
83. Bernstein IL. Cromolyn sodium in the treatment of asthma. Changing concepts. *J Allergy Clin Immunol* 1981;68:247-53.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.